

Electronic Version 1.1

Stylesheet Version v1.1.1

Description

Inhibition of SHIP to Enhance Stem Cell Harvest and Transplantation

BACKGROUND ART

[0001] Statement of Government Interest: The invention described herein may be manufactured and used by the Government of the United States of America for governmental purposes without payment of any royalties thereon.

[0002] Reference to Related Applications: This invention is based on priority document U.S. Provisional Application titled, "Inhibition of SHIP to Enhance Stem Cell Harvest and Transplantation," Application Serial No. 60/320,233 filed on May 28, 2003. This application claims the benefit under 35 U.S.C. §120, as a continuation-in-part (CIP) of the co-pending United States application listed below, and insofar as the subject matter of each of the claims of this application is not disclosed in the manner provided by the first paragraph of 35 U.S.C. §120, I acknowledge the duty to disclose material information as defined in 37 C.F.R. §1.56(a), which occurred between the filing date of this application and the filing date of Application Serial No. 09/955,174, Filed September 19, 2001.

TECHNICAL FIELD

[0003] Traditional treatment for leukemia has consisted of allogeneic transplants of bone marrow (BMT) where the recipient receives tissue grafts from a healthy donor. Although allogeneic BMT has met with some success, rejection of the graft by the recipient as well as the immune response of the grafted tissue attacking the host, known as Graft-Versus-Host-Disease (GVHD), occurs in many cases. GVHD is a significant cause of mortality in allogeneic bone marrow transplant (BMT) procedures. Similarly, attempts to repress the recipient's immune system prior to the transplant leave an already weakened patient susceptible to potentially fatal post-transplant procedures.

[0004] The SH2-domain-containing Inositol 5-Phosphatase (SHIP) is a hematopoietic cell-specific protein that is activated by various cytokines and growth factors. Numerous studies using SHIP deficient mice have shown that SHIP is a negative regulator of hematopoietic cell propagation, function and survival. In essence, SHIP helps control the reproduction of immune cells in the blood known as natural killers (NK) cells. When SHIP is downregulated, or repressed, the immune system of test SHIP-deficient mice were less likely to attack and reject transplanted bone marrow from mismatched, allogeneic donors. The reverse is also true, where the immune system cells present in the transplanted tissue were less likely to attack the new host.

[0005] Down regulation of SHIP in allogeneic BMT recipients prior to transplant has shown a decrease in the number of mice suffering from GVHD. However, allogeneic methods have many variables which affect its

success. In addition to deficiencies already listed, i.e. GVHD and rejection by the host, allogeneic methods are complicated by the ability to find suitable donors and the potentially lethal complications resulting from graft rejection and GVHD.

[0006] For these reasons it is preferable to treat patients with autologous transplant methods, where the patient's own stem cells, depleted of leukemic or tumor cells (including leukemic stem cells clones or tumor cells), are encouraged to grow or are removed, cultured, and re-transplanted into the original host. This method provides the benefits of eliminating the chance of host rejection or GVHD since the cells are not foreign to the body. Currently, these hematopoietic stem cell (HSC) therapies fail with significant frequency. Many times the procedure is infeasible due to the inability to acquire viable HSC from the patient for transplant. Organ-specific stem cell transplants (e.g. neural stem cells, mesenchymal stem cells) are not yet widely used clinically due to the inability to acquire these cells or due to their inability to home efficiently and engraft after transplantation.

[0007] Therefore, what is needed is an efficacious therapy that provides a long-term benefit in a wide variety of genetic, oncologic and infectious diseases in the emerging field of stem cell transplantation.

MODE FOR THE INVENTION

[0008] The instant invention teaches the inhibition of SHIP expression, or function, for the increased efficacy of autologous transplants. In addition to

cancer-related therapies BM stem cells are currently thought to be an effective therapy for cardiac repair. Other tissue or organ-specific stem cells, such as neural stem cells, might also be used to treat neurological diseases that occur due to the loss of neurons in the central nervous system.

[0009] In another embodiment, interference with SHIP function can be used to temporarily expand the hematopoietic stem cell compartment to assist with leukapheresis, to promote hematopoietic recovery after myeloablation treatments, to deplete target stem cell clones (such a leukemic stem cell clones and other tumor stem cell types), and to deplete, or damage, the repopulating ability of the endogenous hematopoietic stem cell pool in order to allow transplanted hematopoietic stem cells to better home and engraft and to increase mesenchymal stem cells number in vivo and ex vivo.

[0010] The method of increasing the yield of stem cells in a patient, in vivo, for autologous transplantation, comprises the steps of administering an effective amount of SHIP inhibitor to the patient and harvesting the stem cells from the patient for autologous transplantation. In one embodiment of this method, the SHIP inhibitor is selected from the group consisting of RNA interference compounds, antisense oligonucleotides, ribozymes, DNazymes, nucleic acid modifiers, PNAs, nonstandard nucleic acids, aptamers, decoys, oligonucleotide based gene regulation, substrate mimics, molecular inhibitors and dominant/negative mutants. The stems cells harvested for transplantation are selected from the group consisting

of hematopoietic stem cells, mammary stem cells, mesenchymal and other organ specific stem cells.

[0011] In another embodiment, the present invention provides a method of increasing the yield of stem cells from a patient, ex vivo, for autologous transplantation, comprising the steps of harvesting target stem cells from the patient and contacting the target stem cells with SHIP inhibitor. The SHIP inhibitor is selected from the group consisting of RNA interference compounds, antisense oligonucleotides, ribozymes, DNAzymes, nucleic acid modifiers, PNAs, nonstandard nucleic acids, aptamers, decoys, oligonucleotide based gene regulation, substrate mimics, molecular inhibitors and dominant/negative mutants. Stems cells harvested for transplantation are selected from the group consisting of hematopoietic stem cells, mammary stem cells, mesenchymal and other organ specific stem cells.

[0012] In yet another embodiment, the present invention provides a non-invasive method for harvesting stem cells from solid organ systems such as the CNS by mobilizing these cells to the blood, comprising the steps of administering SHIP inhibitor to a patient and then harvesting the stem cells from the volume of blood by leukopheresis. The stem cells can be either hematopoietic stem cells (HSC) or non hematopoietic (e.g., CNS, mesenchymal or other organ-specific stem cells).

[0013] In yet another embodiment the present invention provides a non-invasive method of promoting recovery of a stem cell population in a patient comprising the step of administering SHIP inhibitor to the patient, possibly

a patient recovering from myeloablation therapy. The administration of SHIP inhibitor is conducted over a relatively short period, such as approximately one (1) and two (2) weeks. As mentioned, the target stem cell population can be either hematopoietic or non-hematopoietic stem cells.

[0014] In another embodiment, the present invention provides a method of reducing the population of target cells comprising the step of administering an effective amount of SHIP inhibitor to a patient. The administration of SHIP inhibitor is conducted over a more prolonged period, such as approximately nine (9) and twelve (12) weeks and can be used in conjunction with chemotherapeutic agents.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] For a fuller understanding of the nature and objects of the invention, reference should be made to the following detailed description, taken in connection with the accompanying drawings, in which:

[0016] FIG. 1 illustrates how Hematopoietic stem cell (HSC) numbers are increased significantly in SHIP-deficient mice.

[0017] FIG. 2 illustrates how SHIP deficiency increases the number of HSC that are actively cycling.

[0018] FIG. 3 illustrates how mammary stem cells (MSC) are increased significantly in SHIP-deficient mice.

[0019] FIG. 4 illustrates a method to render primary stem cells SHIP deficient.

[0020] Fig. 6 (A) shows flow cytometry analysis of SHIP deficient bone marrow.
(B) Student T test results of the numbers of HSC.

[0021] Fig. 7 represents results from statistical analysis of the numbers
(relative/absolute) of HSC in varying hematopoietic organs.

[0022] Fig. 8 shows the level of reconstitution after a direct competition assay.

[0023] Fig. 9 shows the results of a competitive repopulation assay

[0024] Fig. 10 is a cell cycle analysis on bone marrow from SHIP deficient and
wild type mice.

[0025] Fig. 11 shows that SHIP deficiency alters the size of the mammary stem
cell compartment.

[0026] Fig. 12 shows that SHIP deficiency leads to an increase in mesenchymal
stem cell numbers.

[0027] Fig. 13 is an analysis of SHIP^{+/-} and SHIP^{-/-} Mesenchymal stem cells.

BEST MODE FOR CARRYING OUT THE INVENTION

[0028] The present invention relates to the novel and unexpected finding that
SHIP activity has a crucial role in regulating the number of hematopoietic,
mammary and mesenchymal stem cells. Specifically, the present invention
comprises methods for the increased efficacy of autologous stem cell
transplants.

[0029] Terms: With regard to present disclosure above and below, the following
terms are to be understood as follows.

[0030] SHIP – as used herein refers to the SH2-containing inositol-5-phosphatase-1. SHIP contains an amino-terminal src-homology domain (SH2), a 5'phosphoinositol phosphate domain, two phosphotyrosine binding consensus sequences, and a proline-rich region located in the carboxyl terminus.

[0031] Inhibition of SHIP function – as used herein refers to any genetic, or non-genetic, means for inhibiting SHIP function or expression. The term is meant to comprise any method of achieving this result known in the art. The term includes all methods known, as well as any substance which inhibits SHIP function or expression. Examples include, but are not limited to, RNA interference, antisense oligonucleotides, ribozymes, DNazymes, nucleic acid modifiers, PNAs, nonstandard nucleic acids, aptamers, decoys, oligonucleotide based gene regulation, substrate mimics, molecular inhibitors and dominant/negative mutants.

[0032] Ex Vivo- as used herein refers to transfecting, contacting, or administering a substance to a cell outside of the body. As an example, a cell which shows expression of SHIP is removed from the host. The cell is then transfected, contacted, or administered with a substance that either inhibits SHIP expression or function. The cell is then either transplanted into the host, or allowed to propagate a new population of SHIP deficient cells, in the case of inhibition of SHIP expression, or cells in which SHIP function has been silenced (function).

[0033] Administering or Contacting – as used herein refers to the process of delivering to a cell, ex vivo, or a host, in vivo, a therapeutic substance, or a

combination of several therapeutic substances. The process can include any method known in the art and is dependent on the type of substance or substances administered. Possible methods include, but are not limited to, parenteral (i.e. subcutaneously, intravenously, intramuscularly, intra-arterially, and direct injection into a tissue or organ), mucosal (i.e. intranasally), pulmonary (i.e. via inhalation), topical, via catheter (i.e. iontopheretically) or orally. Administration is usually achieved via a pharmaceutically acceptable carrier.

[0034] Gene Therapy – as used herein refers to a therapeutic method of delivery wherein the therapeutic agent is a polynucleotide capable of accomplishing a therapeutic goal when expressed as a transcript, an interfering RNA, an anti-sense polynucleotide or a polypeptide in the host or cell.

[0035] Therapeutically Effective Amount – as used herein is that amount of a substance necessary to achieve a desired therapeutic result. For example, if the therapeutic result desired is the enhanced yield of stem cells, the therapeutically effective amount is that amount that facilitates, or achieves, an increase in the total number of stem cells in a given population. The therapeutically effective amount can be a dosage administered in at least one amount and can include an administration protocol spanning several days or weeks.

[0036] Genetic Assembly – as used herein refers to any construct which is capable of modulating the expression of a target sequence or gene. Genetic assemblies can include transcriptional promoter/enhancers, locus

defining elements, or any element that controls gene expression by any means (i.e. RNAi, alternate splicing, nuclear RNA export, post-translational modification of intermediaries or proteins). The genetic assembly optimally includes a sequence which is operably linked to the target sequence or gene, when expressed, and acts as a translation initiation sequence.

[0037] Autologous Transplantation

[0038] In one embodiment, the invention provides a method for increasing the yield of stem cells in a patient, in vivo, for autologous transplantation. Such a treatment is valuable for any disease, or genetic impairment, requiring treatment by bone marrow transplant (i.e. cancer, autoimmune deficiencies, HIV/AIDS), blood transplantation (e.g. mobilized stem cells), or stem cell transplant from any tissue source (stem cell transplants may come from non-BM stem cell populations, such as neural stem cells from the CNS, and therefore SHIP inhibition can increase these populations as well, as well as their mobilization to the blood (as it does for hematopoietic stem cells). The non-hematopoietic stem cells can be mobilized to the blood, which would greatly reduce the obstacles faced with collecting cells from bone marrow, brain, or other solid organs). It is known that current hematopoietic stem cell therapies face significant obstacles due to the inability to acquire sufficient numbers of hematopoietic stem cells (HSC) from the patient for transplant. Accordingly, organ-specific stem cell transplants are not currently being done routinely in a clinical setting. In addition to the difficulty in acquiring these cells, efficacious treatments are

hindered by the cells inability to efficiently home and engraft after infusion into circulation.

[0039] Therefore, in one embodiment, a substance that inhibits SHIP is administered to the patient prior to harvesting of the target stem cells. After treatment has been administered for a therapeutically effective time, stem cells are collected and transplanted back into the same host. This has the advantage of avoiding both rejection by the host and Graft Versus Host Disease, wherein the immune response of the grafted tissue attacks the host. Administration of the substance can take the form of genetic assembly or a pharmaceutical agent. These may include nucleic acids, or any substance, that inhibits SHIP function or expression, or leads to the expression of any antagonist of SHIP function. The genetic assembly can be linked to a promoter and other signals directing expression of a protein SHIP antagonist. Antisense oligonucleotides are also viable assemblies. Other possible genetic assemblies include, but are not limited to, RNA interference molecules (sRNA), enzymatic inhibitors, ribozymes, DNazymes, nucleic acid modifiers, dominant/negative mutants, PNAs (or other nonstandard nucleic acids), aptamers, SHIP specific decoys and oligo based gene regulators.

[0040] In another embodiment, the efficiency of autologous tissue transplantation is greatly enhanced by removal of the threat of rejection or GVHD in a patient, ex vivo, through inhibition of SHIP function. Here, stem cells are collected prior to inhibition of SHIP activity. After the cells are collected they are contacted with an effective amount of a substance which inhibits

SHIP activity. The cells are then allowed to propagate to increase their number in the presence of the SHIP inhibitor. When a therapeutically effective population of cells is achieved the cells are transplanted back to the host.

[0041] The number of hematopoietic stem cells (HSC), mammary stem cells (MSC) and mesenchymal stem cells (MeSC) is increased in vivo in SHIP-deficient mice. Thus, inhibition of SHIP expression or activity prior to harvest, or ex vivo after harvesting, allows greater numbers of organ-specific stem cells to be used for transplant and thus ensures that an adequate dose of stem cells is available for transplantation or gene modification. In addition to this quantitative effect, SHIP deficiency increases the proportion of actively dividing stem cells and thus SHIP-deficiency also has a beneficial qualitative effect on stem cell function. SHIP-deficiency also enhances the homing ability of stem cells to the tissue sites that are necessary for their engraftment (e.g. SCF, CXCR4). SHIP is known to increase the migration of differentiated hematopoietic cells to SCF and CXCR4, therefore SHIP inhibition enhances tissue homing of stem cells after their injection or transplantation.

[0042] Particularly, the increased cycling capacity of SHIP-deficient HSC enhances their ability to repopulate a damaged organ and their ability to be transduced by various gene delivery systems. Other features of stem cells are also enhanced by SHIP deficiency, such as homing to the damaged organ or cytokine-stimulated mobilization to the blood for stem cell collection. In a similar manner, inhibition of SHIP expression or activity

can also be used to increase the number of organ-specific stem cells obtained from differentiation of pluripotent stem cells (ES cells). Thus, the development of a method to specifically inhibit SHIP expression or activity is a means to enhance stem cell harvesting, gene modification, homing and collection from the peripheral blood or lymph.

[0043] Example 1 – Effects of in vivo SHIP deficiency on HSC

[0044] Fig. 1 shows that hematopoietic stem cell numbers are increased significantly in SHIP-deficient mice. However, the number of multipotent progenitor cells remains unaffected. Fig.1 (A) shows FACS analysis of SKTL HSC cells and KL stem/progenitors in SHIP^{-/-}, SHIP^{DIP/DIP} and their WT littermates. Fig. 1(B) illustrates absolute and relative numbers of KL stem/progenitors and (C) SKTL HSC in SHIP^{-/-}, SHIP^{DIP/DIP} and their WT littermates. The values determined for SHIP^{-/-} and SHIP^{DIP/DIP} mice that are significantly different from their WT counterparts are indicated by the following symbols: *, $p < 0.05$ and **, $p < 0.0001$. Bone marrow cells were obtained by flushing two intact femurs and two tibias from each mouse. Cells were stained and then analyzed using a FACS Vantage (Becton Dickinson) for the presence of HSC as defined by the Sca1⁺Thy1.2⁺ckit⁺Lin⁻ (SKTL) HSC phenotype. There was no significant difference observed in the absolute or relative number KL stem/progenitor cells present in SHIP mutant strains and their WT counterparts. Note that the background of SHIP^{-/-} mice is essentially C57BL6/J while the SHIP^{DIP/DIP} are F2 (129Sv C57BL6/J). Thus, the SHIP mutation impacts HSC frequency despite differences in genetic

background, further attesting to its role in regulating HSC numbers.

[0045] Example 2 – Proliferation of HSC in the presence of SHIP deficiency

[0046] Fig. 2 shows that a greater proportion of SHIP-deficient HSC are actively proliferating. BrdU staining in Sca+Thy1+kit+Lin- HSC from a representative SHIP^{+/+} (A) and SHIP^{-/-} (B) mouse. (C) Mean BrdU incorporation in Sca+Thy1+kit+Lin- HSC following a 9 day BrdU pulse of SHIP^{-/-} and SHIP^{+/+} mice (n=2 per genotype). The percentage of HSC positive for BrdU incorporation was determined by comparison with an isotype control Ab conjugated to PE (Pharmingen). Only HSC with BrdU staining above that seen in the isotype control were considered positive for BrdU incorporation and this fraction was used to determine the percent BrdU positive HSC. Mice were placed on drinking water containing bromodeoxyuridine (BrdU) at 1mg/ml for 9 days. The mice were then sacrificed and bone marrow cells were isolated from intact tibia and femurs. The BM cells were initially stained with the Lin panel-FITC, c-Kit-APC, Sca1-PE-Cy7 and Thy1.2-Cychrome (eBioscience) antibodies. Following cell permeabilization, the samples were stained with anti-BrdU-PE (BD Pharmingen) and analyzed on a FACS Vantage/DIVA.

[0047] Example 3 - Effects of in vivo SHIP deficiency on MSC

[0048]

Fig. 3. depicts mammary gland stem cells (MSC) as identified by the SP phenotype (Side Population) based on exclusion of Hoechst dye.

Mammary glands were prepared from adult SHIP^{+/+} and SHIP^{-/-} littermates and a single cell suspension was prepared. Viable cells were

analyzed for exclusion of the Hoechst dye and 7AAD. The percentage of MSC present in the mammary gland of each genotype is indicated. Note that there is almost a five-fold increase in the frequency of MSC present in the SHIP^{-/-} mammary glands.

[0049] Example 4 – Example of predictable SHIP inhibition in viv

[0050] Fig. 4. displays one method to render primary stem cells SHIP deficient by RNA interference. Embryonic stem cells that express the SHIP gene were transfected with an irrelevant shRNA vector (Lane 3) or with two different SHIP-specific shRNA vectors (Lanes 4 and 5). The cells were then lysed and equal quantities of whole cell extracts were blotted with either anti-SHIP (Panel A) or anti-Actin (Panel B). Lane 2 shows untreated ES cells. Lane 1 shows untreated RAW264.7 cells that express the SH2 containing SHIP p135 and p145 isoforms. Panel A shows significant reduction of SHIP expression in primary ES cells after transfection of SHIP-specific shRNA vectors in the absence of selection. Please note that these vectors will also interfere with the larger SH2 - containing isoforms expressed in differentiated hematopoietic cells.

[0051] Included herein are thirteen (13) SHIP1 siRNA target sequences (Seq. ID. Nos. 1 – 13). These sequences are predicted to have strong specificity and strong knockdown against the human SHIP1 cDNA sequence. The cDNA is about 4800+ nt (Seq. ID. No. 14). Accordingly other target sequences are available, however, those listed (13) have a strong predictability factor in vivo.

[0052] Long Term Repopulation Modulation: In addition to the present inventions unexpected findings with regard to the enhancement of stem populations in a SHIP deficient environment, the present invention also relates to the novel and unexpected finding that SHIP deficient cells experience difficulty with regard to long-term repopulation of the blood system in comparison to wild-type hematopoietic stem cells.

[0053] Therefore, in another embodiment, interference with SHIP function can be used to temporarily expand the hematopoietic stem cell compartment to assist with leukapheresis, to promote hematopoietic recovery after myeloablation treatments, to deplete target stem cell clones (such a leukemic clones and other tumor stem cell types), and to deplete, or damage, the repopulating ability of the endogenous hematopoietic stem cell pool in order to allow transplanted hematopoietic stem cells to better home and engraft and to increase mesenchymal and stem cells number in vivo and ex vivo.

[0054] Figs. 2 and 10 show that the frequency of HSC that are actively cycling is increased and that HSC numbers are increased in the blood (Fig. 5). These findings show SHIP inhibition could be used to increase the number of cycling (i.e. actively-dividing) stem cells and thus their mobilization to the peripheral blood. The cycling of other stem cell types (mesenchymal, mammary, neural or organ-specific stem cells) may also be increased by SHIP-deficiency and thus non-hematopoietic stem cells may also be mobilized from their tissue of origin to the peripheral blood (thus, enabling non-invasive harvesting methods for non-hematopoietic stem cells by

leukapheresis). Accordingly, temporary period of SHIP increases the cycling rate of endogenous stem cell populations and their mobilization to the blood for harvest. However, this period of SHIP inhibition should only be temporary (~1-2 weeks), as prolonged SHIP-deficiency (6-10 weeks) likely causes aging of endogenous stem cells as seen by the reduced ability of HSC in adult SHIP-deficient mice to repopulate the blood or tissue of interest (see Figs 8,9).

[0055] Alternatively, endogenous stem cells may not need to be harvested for subsequent return to the patient. The patient could be treated for a temporary period with SHIP inhibitors to increase the cycling of HSC to allow more rapid recovery of key blood cell populations (e.g. platelets, granulocytes, erythroid cells) after myeloablation. The same principle could also be applied to other stem cell populations in diseases where damage to tissues can be repaired by increasing the cycling and repopulation capacity of the stem cells for that tissue.

[0056] The ability of normal HSC to contribute to normal blood cell production is damaged by prolonged SHIP deficiency (9-12 weeks). This may occur by increasing the division rate of these cells and therefore prematurely “aging” these cells. Therefore, prolonged SHIP deficiency might also be used to burn out or “age” leukemic or tumor stem cell clones that must replenish cells of the leukemia or tumor. SHIP inhibitors might be combined with chemotherapeutic or other anti-cancer agents.

[0057] Treatment with SHIP inhibitors could also be used to age endogenous stem cell populations so that stem cell transplants (autologous or

allogeneic) could compete more effectively for the tissues niches that support stem cell engraftment.

[0058] Example 5 – SHIP deficiency increases the absolute number of HSC in vivo, increases their relative proportion and mobilizes HSC to the peripheral blood.

[0059] Fig. 5 (A) shows bone marrow (BM), spleen, and peripheral blood (PB) from SHIP^{-/-} H2B mice were analyzed for the presence of hematopoietic stem cells (HSC;KTLS:Lin-cKit+Sca1+ Thy1+) and early progenitor cells (Lin-cKit+) and late progenitor cells (LinlowcKit+) by flow cytometry analysis. B) Bone marrow from SHIPDIP/DIP was analyzed for HSC and progenitor cells as mentioned for BM H2B. C) Statistical analysis showing the relative and absolute numbers of HSC and progenitor cells different hematopoietic organs. The statistical analysis was performed using the unpaired Student T test. Results were considered significant (indicated by an *) when p values were under 0.05.

[0060] Example 6 – Method and results of SHIP ablation during adulthood (in vivo)

[0061] In Fig. 6 HSC; KTLS (Lin-cKit+Sca1+Thy1+) cells are increased in mice in which SHIP was ablated during adulthood. (A) Bone marrow (BM) from Mx-CRE SHIP^{-/-} and SHIP^{+/-} mice were analyzed for the presence of hematopoietic stem cells (HSC; Lin-cKit+Sca1+Thy1+) by flow cytometry analysis. Mx-CRE SHIP^{-/-} mice are created by injecting MxCRE SHIPFlox^{-/-} mice with polyIC, which induces the production of interferon.

Interferon acts on the Mx promoter to cause the transcription of CRE. CRE will then cause recombination of the flox site, leading to SHIP deletion.

This model allow use to study the effect of SHIP removal during adulthood. B) Statistical analysis showing the relative numbers of HSC from Mx-CRE SHIP^{-/-} and SHIP^{-/flox}.

[0062] The statistical analysis was performed using the unpaired Student t test in Prism 4. Results were considered significant when $p < 0.005$. * $p < 0.05$.

[0063] In Fig. 7 the same group that defined the KTLS phenotype, found another method to isolate immunophenotype to isolate HSC, which is Lin-Flk2-cKit+Sca1+ (LFKS). LFKS cells are also increased in the SHIP^{-/-} mice. (A) Bone marrow (BM), spleen, and peripheral blood (PB) from SHIP^{-/-} H2B mice were analyzed for the presence of hematopoietic stem cells following a different immunophenotype (HSC; Lin-Flk2-cKit+Sca1+) by flow cytometry analysis, on FACS Calibur. B) Statistical analysis showing the relative numbers of HSC in the different hematopoietic organs. C) Statistical analysis showing the absolute numbers of HSC in the different hematopoietic organs. The statistical analysis was performed using the unpaired Student T test. Results were considered significant (indicated by an *) when p values were under 0.05.

[0064] Fig. 8 shows the level of reconstitution after direct competition assay.

Direct competition assay was performed by sorting KTLS (Lin-cKit+Sca1+Thy1+) cells from SHIP^{-/-} Ly5.2 mice and from WT Ly5.1 mice. 200 SHIP^{-/-} Ly5.2 and 200 WT Ly5.1 KTLS were then injected into the same animal with 40 000 Sca1⁻ supporting cells (Ly5.1/Ly5.2). The

recipient mice were Ly5.1/Ly5.2, which allow us to identify which cells in the immune system comes from the recipient, WT donor or SHIP^{-/-} donor. The recipient mice had undergone total body irradiation (dose: 600 rads and 2 hours later 400 rads) 2 hours before being transplanted with the test cells. After transplantation the mice were given antibiotic water to prevent opportunistic pathogen. At different time point after transplantation, the mice were tested for reconstitution of the hematopoietic compartment. A) Global reconstitution 4 weeks and 4 month after transplantation. The results show no significant difference 4 weeks after transplantation between the WT and the SHIP^{-/-} HSC ability to reconstitute the hematopoietic system.

[0065] Four (4) months after transplantation, the proportion of hematopoietic cells derived from the WT KTLS is significantly higher than the one derived from SHIP^{-/-} KTLS cells. B) shows the level of reconstitution, 4 months after transplantation, in 3 different hematopoietic lineage to show that the cells are pluripotent. Again the level of reconstitution from the WTKTLS is significantly higher than from SHIP^{-/-} KTLS. This is the result of 11 mice done in two different experiments. Statistical significance was established using Prism 4 software,

[0066] unpaired student t test. + $p > 0.05$, * $p < 0.01$, ** $p < 0.005$, *** $p < 0.05$.

[0067] Example 7 - Long term competition effects of SHIP deficiency

[0068] Fig. 9 is a competitive repopulation assay showing that SHIP^{-/-} bone marrow has less CRU than WT littermates BM. Mice transplanted: Ly5.1

C57Bl6 mice that were irradiated with one-single dose of 950 rads. The tested cells were Ly5.2 cells from either SHIP^{-/-} mice or WT mice and the competing cells were Ly5.1.

[0069] (A) Level of reconstitution of the hematopoietic system by the tested donor BM. WT BM level of reconstitution is significantly higher than for SHIP^{-/-} BM. Student t test $p < 0.0001$. (B) Number of competitive repopulation unit (CRU) was established following a method developed by D.E. Harrison, where donor CRU = $(10 \times \% \text{donor}) / (100 - \% \text{donor})$. 10 is the number of CRU present in the competing BM (Ly5.1). WT BM has a significantly higher number of CRU compared to SHIP^{-/-} mice. Unpaired t test $p = 0.0005$.

[0070] Fig. 10 demonstrates that cell cycle analysis on BM from SHIP deficient and WT mice revealed that a greater proportion of SHIP^{-/-} HSC in cell cycle. Bar graph representing the proportion of Lin-cKit+Sca1⁺ cells that are G0 (resting) or in S/G2 phase (dividing). This graph includes results from experiment performed using SHIP^{-/-} mice on a C57Bl6 background and SHIPDIP/DIP on a 129SvJ background with respective WT counterparts.

[0071] Example 8 – Effects of SHIP deficiency in non-HSC

[0072]

Fig. 11 shows that mammary stem cells were analyzed from both H2B and H2D SHIP mice and from Δ E IP (Rock) mice. Cells were isolated from the third and fourth mammary glands of 6-8 week old female mice. The glands were made into a single cell suspension through physical processing and

enzymatic digestion. The mammary gland single cell suspension was examined for MaSC using the lineage negative gate of CD45-, and two positive gates SP+ and Sca+.

[0073] Fig. 12 MeSC were analyzed from SHIP+/+ and SHIP-/- mice on a C57Bl6/J background. Femurs and tibiae were collected from three sets of mice. The muscle, cartilage, and marrow were removed. The bone was crushed with a mortar and pestle and the fragments were digested with collagenase. Cells were plated with MeSC isolation media in triplicate at equal density. They were allowed to attach for 24 hours at the conclusion of which the non-adherent cells were washed away and the adherent cells were counted. A students t-test was performed and the P-value < 0.0001.

[0074] Fig. 13 represents the analysis of SHIP+/- and SHIP-/- MeSC. Bar graphs represent the number of MeSC per 1million input whole bone marrow cells plated at time zero. Cells were counted, after lifting with trypsin, at 24 hours and 4 weeks post time zero using a hemacytometer. Total cell numbers decreased at 4 weeks as the cultures became a more homogenous population, though the difference that exists between the SHIP+/+ and SHIP-/- cell counts at each time point increased from a factor of 3.2 for the WT to 11.7 for the null which has a p value of $p < 0.003$ indicating that this increase is significant. * $p < 0.0001$ ** $p < 0.01$ by a two-tailed students t-test.

[0075] It will be seen that the objects set forth above, and those made apparent from the foregoing description, are efficiently attained and since certain changes may be made in the above construction without departing from

the scope of the invention, it is intended that all matters contained in the foregoing description or shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

[0076] It is also to be understood that the following claims are intended to cover all of the generic and specific features of the invention herein described, and all statements of the scope of the invention which, as a matter of language, might be the to fall therebetween. Now that the invention has been described,

SEQUENCE LISTING

[0077] <110> University of South Florida

[0078] <120> INHIBITION OF SHIP TO ENHANCE STEM CELL HARVEST AND

[0079] TRANSPLANTATION

[0080] <130> 1372.160PRC

[0081] <160> 14

[0082] <170> PatentIn version 3.2

[0083] <210> 1

[0084] <211> 19

[0085] <212> RNA

[0086] <213> Artificial Sequence

[0087] <220>

[0088] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0089] <400> 1

[0090] gcctgtgtgc atccattga 19

[0091] <210> 2

[0092] <211> 19

[0093] <212> RNA

[0094] <213> Artificial Sequence

[0095] <220>

[0096] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0097] <400> 2

[0098] ataagttggt gatcttggt 19

[0099] <210> 3

[0100] <211> 19

[0101] <212> RNA

[0102] <213> Artificial Sequence

[0103] <220>

[0104] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0105] <400> 3

[0106] gccacatctg tactgacaa 19

[0107] <210> 4

[0108] <211> 19

[0109] <212> RNA

[0110] <213> Artificial Sequence

[0111] <220>

[0112] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0113] <400> 4

[0114] agacagggcat tgcaaacac 19

[0115] <210> 5

[0116] <211> 19

[0117] <212> RNA

[0118] <213> Artificial Sequence

[0119] <220>

[0120] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0121] <400> 5

[0122] acatcactca ccgcttcac 19

[0123] <210> 6

[0124] <211> 19

[0125] <212> RNA

[0126] <213> Artificial Sequence

[0127] <220>

[0128] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0129] <400> 6

[0130] tcttaactac cgtgtggat 19

[0131] <210> 7

[0132] <211> 19

[0133] <212> RNA

[0134] <213> Artificial Sequence

[0135] <220>

[0136] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0137] <400> 7

[0138] aatacgccta caccaagca 19

[0139] <210> 8

[0140] <211> 19

[0141] <212> RNA

[0142] <213> Artificial Sequence

[0143] <220>

[0144] <223> SHIP1 siRNA target sequences. Predicted to have good specificity

[0145] and good knockdown against the human SHIP1 cDNA sequence.

[0146] <400> 8

[0147] gtaccagcga catcatgac 19

[0148] <210> 9

[0149] <211> 19

[0150] <212> RNA

[0151] <213> Artificial Sequence

[0152] <220>

[0153] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0154] <400> 9

[0155] gcgacatcat gacgagtga 19

[0156] <210> 10

[0157] <211> 19

[0158] <212> RNA

[0159] <213> Artificial Sequence

[0160] <220>

[0161] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0162] <400> 10

[0163] aggacagatt gagtttctc 19

[0164] <210> 11

[0165] <211> 19

[0166] <212> RNA

[0167] <213> Artificial Sequence

[0168] <220>

[0169] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0170] <400> 11

[0171] ggtgctatgc cacattgaa 19

[0172] <210> 12

[0173] <211> 19

[0174] <212> RNA

[0175] <213> Artificial Sequence

[0176] <220>

[0177] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0178] <400> 12

[0179] gtttggtgag actcttcca 19

[0180] <210> 13

[0181] <211> 19

[0182] <212> RNA

[0183] <213> Artificial Sequence

[0184] <220>

[0185] <223> SHIP1 siRNA target sequences. Predicted to have good specificity and good knockdown against the human SHIP1 cDNA sequence.

[0186] <400> 13

[0187] agacggagcg tgatgaatc 19

[0188] <210> 14

[0189] <211> 4870

[0190] <212> DNA

[0191] <213> Human

[0192] <400> 14

[0193] gtggaggggc ctccgctccc ctcggtggtg tgtgggtcct gggggtgcct gccggcccag 60

[0194] ccgaggaggc ccacgccac catggtcccc tgctggaacc atggcaacat caccgctcc
120

[0195] aaggcggagg agctgcttc caggacaggc aagggcacga gcttcctcgt gcgtgccagc
180

[0196] gagtccatct cccgggcata cgcgctctgc gtgctgtatc ggaattgcgt ttacacttac 240

[0197] agaattctgc ccaatgaaga tgataaatc actgttcagg catccgaagg cgtctccatg 300

[0198] aggttcttca ccaagctgga ccagctcatc gagttttaca agaaggaaaa catggggctg 360

[0199] gtgacccatc tgcaataccc tgtgccgctg gaggaagagg acacaggcga cgaccctgag
420

[0200] gaggacacag tagaaagtgt cgtgtctcca cccgagctgc cccaagaaa catcccgctg
480

[0201] actgccagct cctgtgaggc caaggagggt ccttttcaa acgagaatcc ccgagcgacc 540

[0202] gagaccagcc ggccgagcct ctccgagaca ttgtccagc gactgcaaag catggacacc
600

[0203] agtgggcttc cagaagagca tcttaaggcc atccaagatt atttaagcac tcagctcgcc 660

[0204] caggactctg aatttgtaa gacaggggtc agcagtcttc ctcacctgaa gaaactgacc 720

[0205] aactgtctct gcaaggagct ctatggagaa gtcacccgga ccctcccatc cctggagtct 780

[0206] ctgcagaggt tattgacca gcagctctcc ccgggcctcc gtccacgtcc tcaggttctt 840

[0207] ggtgaggcca atcccatcaa catggtgtcc aagctcagcc aactgacaag cctgttgtca 900

[0208] ccattgaag acaagggtcaa ggccttgctg cacgagggtc ctgagtctcc gcaccggccc 960

[0209] tcccttatcc ctccagtcac ctttgaggtg aaggcagagt ctctggggat tcctcagaaa 1020

[0210] atgcagctca aagtcgacgt tgagtctggg aaactgatca ttaagaagtc caaggatggt 1080

[0211] tctgaggaca agttctacag ccacaagaaa atcctgcagc tcattaagtc acagaaattt 1140

[0212] ctgaataagt tggatgctt ggtggaaaca gagaaggaga agatcctgcg gaaggaatat
1200

[0213] gtttttgctg actcaaaaa gagagaaggc ttctgccagc tcctgcagca gatgaagaac 1260

[0214] aagcactcag agcagccgga gcccgacatg atcaccatct tcacggcac ctggaacatg
1320

[0215] ggtaacgccc cccctcccaa gaagatcacg tcctggttc tctccaaggg gcagggaaag
1380

[0216] acgcgggacg actctgcgga ctacatcccc catgacattt acgtgatcgg cacccaagag
1440

[0217] gaccccctga gtgagaagga gtggctggag atcctcaaac actccctgca agaaatcacc
1500

[0218] agtgtgactt taaaacagt cgccatccac acgctctgga acatccgcat cgtggtgctg 1560

[0219] gccaaagcctg agcacgagaa ccgcatcagc cacatctgta ctgacaacgt gaagacaggc
1620

[0220] attgcaaaca cactggggaa caaggagacc gtgggggtgt cgttcattt caatggaacc
1680

[0221] tccttagggt tcgtcaacag ccactgact tcaggaagtg aaaagaaact caggcgaaac
1740

[0222] caaaactata tgaacattct ccggttcctg gccctgggcg acaagaagct gagtccctt 1800

[0223] aacatcactc accgcttcac gcacctctc tggttgggg atcttaacta ccgtgtggat 1860

[0224] ctgcctacct gggaggcaga aaccatcatc cagaaaatca agcagcagca gtacgcagac
1920

[0225] ctctgtccc acgaccagct gctcacagag aggaggagc agaaggctt cctacactc
1980

[0226]
gaggaggaag aaatcacgtt tgccccaacc tacggtttg agagactgac tcgggacaaa

2040

[0227] tacgcctaca ccaagcagaa agcgacaggg atgaagtaca acttgccttc ctggtgtgac

2100

[0228] cgagtcctct ggaagtctta tcccctgggtg cacgtgggtgt gtcagtctta tggcagtacc 2160

[0229] agcgacatca tgacgagtga ccacagccct gtctttgcca catttgaggc aggagtcaact 2220

[0230] tcccagtttg tctccaagaa cgggtcccggg actgttgaca gccaaggaca gattgagttt 2280

[0231] ctgaggtgct atgccacatt gaagaccaag tcccagacca aattctacct ggagttccac 2340

[0232] tcgagctgct tggagagttt tgtcaagagt caggaaggag aaaatgaaga aggaagtgag

2400

[0233] ggggagctgg tggtaagtt tggtagact ctccaaagc tgaagcccat tatctctgac 2460

[0234] cctgagtacc tgctagacca gcacatctc atcagcatca agtcctctga cagcgacgaa

2520

[0235] tcctatggcg agggctgcat tgcccttcgg ttagaggcca cagaaacgca gctgcccac 2580

[0236] tacacgcctc tcaccacca tggggagttg acaggccact tccaggggga gatcaagctg

2640

[0237] cagacctctc agggcaagac gagggagaag ctctatgact ttgtgaagac ggagcgtgat

2700

[0238] gaatccagtg ggccaaagac cctgaagagc ctaccagcc acgaccccat gaagcagtgg

2760

[0239] gaagtcacta gcagggcccc tccgtgcagt ggctccagca tcaactgaaat catcaacccc

2820

[0240] aactacatgg gagtggggcc ctttgggccca ccaatgcccc tgcacgtgaa gcagaccttg

2880

[0241] tcccctgacc agcagcccac agcctggagc tacgaccagc cgcccaagga ctccccgctg

2940

[0242] gggccctgca ggggagaaag tcctccgaca cctcccggcc agccgcccc atcacccaag

3000

[0243] aagtttttac cctcaacagc aaaccggggg ctccctccca ggacacagga gtcaaggccc

3060

[0244] agtgacctgg ggaagaacgc aggggacacg ctgcctcagg aggacctgcc gctgacgaag

3120

[0245] cccgagatgt ttgagaaccc cctgtatggg tccctgagtt ccttcataa gcctgctccc 3180

[0246] aggaaggacc aggaatcccc caaatgccg cggaaggaac ccccgccctg cccggaaccc

3240

[0247] ggcatcttgt cgcccagcat cgtgctcacc aaagcccagg aggctgatcg cggcgagggg

3300

[0248] cccggcaagc aggtgcccgc gccccggctg cgctccttca cgtgctcatc ctctgccgag 3360

[0249] ggcagggcgg ccggcgggga caagagccaa ggaagccca agaccccgt

cagctcccag 3420

[0250]

gccccggtgc cggccaagag gcccatcaag ccttcagat cggaatatcaa ccagcagacc

3480

[0251] ccgcccaccc cgacgccgcg gccgccgctg ccagtcaaga gcccggcggt gctgcacctc

3540

[0252] cagcactcca agggccgcga ctaccgcgac aacaccgagc tcccgcata cggaagcac

3600

[0253] cggccggagg aggggccacc agggcctcta ggcaggactg ccatgcagtg aagccctcag

3660

[0254] tgagctgcca ctgagtcggg agcccagagg aacggcgtga agccactgga ccctctccc

3720

[0255] ggacctctg ctggctctc ctgccagct tcctatgcaa ggctttgtgt ttcaggaaa 3780

[0256] gggcctagct tctgtgtggc ccacagagtt cactgcctgt gagacttagc accaagtgt 3840

[0257] gaggctggaa gaaaaacgca caccagacgg gcaacaaaca gtctgggtcc ccagctcgct

3900

[0258] cttggtactt gggaccccag tgccctgttg agggcgccat tctgaagaaa ggaactgcag 3960

[0259] cgccgatttg agggtggaaga tatagataat aataatatta ataataataa tggccacatg 4020

[0260] gatcgaacac tcatgatgtg ccaagtgtg tgctaagtgc ttacgaaca ttcgtcatat 4080

[0261] caggatgacc tcgagagctg aggctctagc cacctaaaac cacgtgccca aaccaccag

4140

[0262] tttaaaacgg tgtgtgttcg gaggggtgaa agcattaaga agcccagtc cctcctggag 4200

[0263] tgaaacaagg gctcggcctt aaggagctga agagtctggg tagcttgtt aggggtacaag 4260

[0264] aagcctgttc tgtccagctt cagtgcacaca agctgcttta gctaaagtcc cgcggggtcc 4320

[0265] ggcatggcta ggctgagagc agggatctac ctggcttctc agttctttgg ttggaaggag 4380

[0266] caggaaatca gctcctattc tccagtggag agatctggcc tcagcttggg ctagagatgc 4440

[0267] caaggcctgt gccagggtcc ctgtgccctc ctcgagggtg gcagccatca ccagccacag
4500

[0268] ttaagccaag ccccccaaca tgtattccat cgtgctggta gaagagtctt tgctgttgct 4560

[0269] cccgaaagcc gtgctctcca tcctggctgc cagggagggt gggcctcttg gttccaggct 4620

[0270] cttgaaatag tgcagccttt tcttctatc tctgtggctt tcaactctgc ttccttggtt 4680

[0271] attaagagaa tagatgggtg atgtctttcc ttatgttgct tttcaacat agcagaatta 4740

[0272] atgttgggag ctaaattcac tgggtgtgtg gaatgcagaa gggaatgcac cccaccttcc 4800

[0273] catgaatgaa gtctgcgtac caataaattg tgccttctcc tcaaaaaaaaa aaaaaaaaaa
4860

[0274] ataaaaaaaa 4870